Three-Pion HBT Correlations in Relativistic Heavy-Ion Collisions from the STAR Experiment

J. Adams[3], C. Adler[11], Z. Ahammed[24], C. Allgower[12], J. Amonett[14], B.D. Anderson[14], M. Anderson[5], D. Arkhipkin[10], G.S. Averichev[9], J. Balewski[12], O. Barannikova[9,24], L.S. Barnby[14], J. Baudot[13], S. Bekele[21], V.V. Belaga[9], R. Bellwied[33], J. Berger[11], H. Bichsel[32], A. Billmeier[33], L.C. Bland[2], C.O. Blyth[3], B.E. Bonner[25], M. Botje[20], A. Boucham[28], A. Brandin[18], A. Bravar[2], R.V. Cadman[1], X.Z. Cai[27], H. Caines[35], M. Calderón de la Barca Sánchez[2], A. Cardenas[24], J. Carroll[15], J. Castillo[15], M. Castro[33], D. Cebra[5], P. Chaloupka[21], S. Chattopadhyay[33], Y. Chen[6], S.P. Chernenko[9], M. Cherney[8], A. Chikanian[35], B. Choi[30], W. Christie[2], J.P. Coffin[13], T.M. Cormier[33], M. Mora Corral[16], J.G. Cramer[32], H.J. Crawford[4], A.A. Derevschikov[23], L. Didenko[2], T. Dietel[11], J.E. Draper[5], V.B. Dunin[9], J.C. Dunlop[35], V. Eckardt[16], L.G. Efimov[9], V. Emelianov[18], J. Engelage[4], G. Eppley[25], B. Erazmus[28], P. Fachini[2], V. Faine[2], J. Faivre[13], R. Fatemi[12], K. Filimonov[15], E. Finch[35], Y. Fisyak[2], D. Flierl[11], K.J. Foley[2], J. Fu[15,34], C.A. Gagliardi[29], N. Gagunashvili[9], J. Gans[35], L. Gaudichet[28], M. Germain[13], F. Geurts[25], V. Ghazikhanian[6], O. Grachov[33], M. Guedon[13], S.M. Guertin[6], E. Gushin[18], T.D. Gutierrez[5], T.J. Hallman[2], D. Hardtke[15], J.W. Harris[35], M. Heinz[35], T.W. Henry[29], S. Heppelmann[22], T. Herston[24], B. Hippolyte[13], A. Hirsch[24], E. Hjort[15], G.W. Hoffmann[30], M. Horsley[35], H.Z. Huang[6], T.J. Humanic[21], G. Igo[6], A. Ishihara[30], P. Jacobs[15], W.W. Jacobs[12], M. Janik[31], I. Johnson[15], P.G. Jones[3], E.G. Judd[4], S. Kabana[35], M. Kaneta[15], M. Kaplan[7], D. Keane[14], J. Kiryluk[6], A. Kisiel[31], J. Klay[15], S.R. Klein[15], A. Klyachko[12], T. Kollegger[11], A.S. Konstantinov[23], M. Kopytine[14], L. Kotchenda[18], A.D. Kovalenko[9], M. Kramer[19], P. Kravtsov[18], K. Krueger[1], C. Kuhn[13], A.I. Kulikov[9], G.J. Kunde[35], C.L. Kunz[7], R.Kh. Kutuev[10], A.A. Kuznetsov[9], M.A.C. Lamont[3], J.M. Landgraf[2], S. Lange[11], C.P. Lansdell[30], B. Lasiuk[35], F. Laue[2], J. Lauret[2], A. Lebedev[2], R. Lednický[9], V.M. Leontiev[23], M.J. LeVine[2], Q. Li[33], S.J. Lindenbaum[19], M.A. Lisa[21], F. Liu[34], L. Liu[34], Z. Liu[34], Q.J. Liu[32], T. Ljubicic[2], W.J. Llope[25], H. Long[6], R.S. Longacre[2], M. Lopez-Noriega[21], W.A. Love[2], T. Ludlam[2], D. Lynn[2], J. Ma[6], Y.G. Ma[27], D. Magestro[21], R. Majka[35], S. Margetis[14], C. Markert[35], L. Martin[28], J. Marx[15], H.S. Matis[15], Yu.A. Matulenko[23], T.S. McShane[8], F. Meissner[15], Yu. Melnick[23], A. Meschanin[23], M. Messer[2], M.L. Miller[35], Z. Milosevich[7], N.G. Minaev[23], J. Mitchell[25], L. Molnar[24], C.F. Moore[30], V. Morozov[15], M.M. de Moura[33], M.G. Munhoz[26], J.M. Nelson[3], P. Nevski[2], V.A. Nikitin[10], L.V. Nogach[23], B. Norman[14], S.B. Nurushev[23], G. Odyniec[15], A. Ogawa[2], V. Okorokov[18], M. Oldenburg[16], D. Olson[15], G. Paic[21], S.U. Pandey[33], Y. Panebratsev[9], S.Y. Panitkin[2], A.I. Pavlinov[33], T. Pawlak[31], V. Perevoztchikov[2], W. Peryt[31], V.A Petrov[10], R. Picha[5], M. Planinic[12], J. Pluta[31], N. Porile[24], J. Porter[2], A.M. Poskanzer[15], E. Potrebenikova[9], D. Prindle[32], C. Pruneau[33], J. Putschke[16], G. Rai[15], G. Rakness[12], O. Ravel[28], R.L. Ray[30], S.V. Razin[9,12], D. Reichhold[24], J.G. Reid[32], G. Renault[28], F. Retiere[15], A. Ridiger[18], H.G. Ritter[15], J.B. Roberts[25], O.V. Rogachevski[9], J.L. Romero[5], A. Rose[33], C. Roy[28], V. Rykov[33], I. Sakrejda[15], S. Salur[35], J. Sandweiss[35], I. Savin[10], J. Schambach[30], R.P. Scharenberg[24], N. Schmitz[16], L.S. Schroeder[15], K. Schweda[15], J. Seger[8], P. Seyboth[16], E. Shahaliev[9], K.E. Shestermanov[23], S.S. Shimanskii[9], F. Simon[16], G. Skoro[9], N. Smirnov[35], R. Snellings[20], P. Sorensen[6], J. Sowinski[12], H.M. Spinka[1], B. Srivastava[24], E.J. Stephenson[12], R. Stock[11], A. Stolpovsky[33], M. Strikhanov[18], B. Stringfellow[24], C. Struck[11], A.A.P. Suaide[33], E. Sugarbaker[21], C. Suire[2], M. Šumbera[21], B. Surrow[2], T.J.M. Symons[15], A. Szanto de Toledo[26], P. Szarwas[31], A. Tai[6], J. Takahashi[26], A.H. Tang[15], D. Thein[6], J.H. Thomas[15], M. Thompson[3], S. Timoshenko[18], M. Tokarev[9], M.B. Tonjes[17], T.A. Trainor[32], S. Trentalange[6], R.E. Tribble[29], V. Trofimov[18], O. Tsai[6], T. Ullrich[2], D.G. Underwood[1], G. Van Buren[2], A.M. Vander Molen[17], A.N. Vasiliev[23], S.E. Vigdor[12], S.A. Voloshin[33], M. Vznuzdaev[18], F. Wang[24], Y. Wang[30], H. Ward[30], J.W. Watson[14], R. Wells[21], G.D. Westfall[17], C. Whitten Jr. [6], H. Wieman[15], R. Willson[21], S.W. Wissink[12], R. Witt[35], J. Wood[6], N. Xu[15], Z. Xu[2], A.E. Yakutin[23], E. Yamamoto[15], J. Yang[6], P. Yepes[25], V.I. Yurevich[9], Y.V. Zanevski[9], I. Zborovský[9], H. Zhang[35], W.M. Zhang[14], R. Zoulkarneev[10], J. Zoulkarneeva[10], A.N. Zubarev[9]

(STAR Collaboration)

[1] *Argonne National Laboratory, Argonne, Illinois 60439*
[2] *Brookhaven National Laboratory, Upton, New York 11973*
[3] *University of Birmingham, Birmingham, United Kingdom*
[4] *University of California, Berkeley, California 94720*
[5] *University of California, Davis, California 95616*

[6] *University of California, Los Angeles, California 90095*
[7] *Carnegie Mellon University, Pittsburgh, Pennsylvania 15213*
[8] *Creighton University, Omaha, Nebraska 68178*
[9] *Laboratory for High Energy (JINR), Dubna, Russia*
[10] *Particle Physics Laboratory (JINR), Dubna, Russia*
[11] *University of Frankfurt, Frankfurt, Germany*
[12] *Indiana University, Bloomington, Indiana 47408*
[13] *Institut de Recherches Subatomiques, Strasbourg, France*
[14] *Kent State University, Kent, Ohio 44242*
[15] *Lawrence Berkeley National Laboratory, Berkeley, California 94720*
[16] *Max-Planck-Institut fuer Physik, Munich, Germany*
[17] *Michigan State University, East Lansing, Michigan 48825*
[18] *Moscow Engineering Physics Institute, Moscow Russia*
[19] *City College of New York, New York City, New York 10031*
[20] *NIKHEF, Amsterdam, The Netherlands*
[21] *Ohio State University, Columbus, Ohio 43210*
[22] *Pennsylvania State University, University Park, Pennsylvania 16802*
[23] *Institute of High Energy Physics, Protvino, Russia*
[24] *Purdue University, West Lafayette, Indiana 47907*
[25] *Rice University, Houston, Texas 77251*
[26] *Universidade de Sao Paulo, Sao Paulo, Brazil*
[27] *Shanghai Institute of Nuclear Research, Shanghai 201800 China*
[28] *SUBATECH, Nantes, France*
[29] *Texas A&M University, College Station, Texas 77843*
[30] *University of Texas, Austin, Texas 78712*
[31] *Warsaw University of Technology, Warsaw, Poland*
[32] *University of Washington, Seattle, Washington 98195*
[33] *Wayne State University, Detroit, Michigan 48201*
[34] *Institute of Particle Physics, CCNU (HZNU), Wuhan, 430079 China*
[35] *Yale University, New Haven, Connecticut 06520*

(Dated: June 20, 2003)

Data from the first physics run at the Relativistic Heavy-Ion Collider at Brookhaven National Laboratory, Au+Au collisions at $\sqrt{s_{NN}} = 130$ GeV, have been analyzed by the STAR Collaboration using three-pion correlations with charged pions to study whether pions are emitted independently at freezeout. We have made a high-statistics measurement of the three-pion correlation function and calculated the normalized three-particle correlator to obtain a quantitative measurement of the degree of chaoticity of the pion source. It is found that the degree of chaoticity seems to increase with increasing particle multiplicity.

PACS numbers: 25.75.Gz, 25.75.Ld

Two-pion Hanbury Brown and Twiss (HBT) interferometry in principle provides a means of extracting the space-time evolution of the pion source produced at kinematic freeze-out in relativistic heavy-ion collisions [1, 2]. An underlying assumption of this method is that pions are produced from a completely chaotic source, i.e. a source in which the hadronized pions are created with random quantum particle production phases. In applications of two-pion HBT the validity of this assumption is usually tested by extracting the "λ-parameter" which in a simple picture is unity for a fully chaotic source and zero for a fully coherent source [1, 2]. However, this parameter also depends on many other factors, such as contamination from other particles in the pion sample, unresolvable contributions from the decay of long-lived resonances and unstable particles (ω, η, η', K^0, Λ, etc.), and inaccurate Coulomb corrections [2].

A better determination of the source chaoticity is possible by using three-particle correlations. Normalizing the three-pion correlation function appropriately by the two-pion correlator, the effects from particle misidentification and decay contributions can be removed [3], thereby isolating possible coherence effects in the particle emission process. The resulting three-pion correlator

r_3 provides the means of extracting the degree of source chaoticity by examining its value in the limit of zero relative momentum. Recent measurements at the CERN SPS from experiments NA44 and WA98 have focused on extracting r_3 from three-pion correlations [4, 5]. While these studies have produced results which are consistent with a chaotic source for Pb+Pb collisions ($\sqrt{s_{NN}} = 17$ GeV), NA44 in particular has shown for S+Pb collisions ($\sqrt{s_{NN}} = 20$ GeV) a result which does not appear to be consistent with the chaotic assumption. All of these prior results suffer from low statistics which limits their significance. We present here using charged pions the first high-statistics heavy-ion study of three-pion correlations, resulting in the first accurate measurement of the degree of chaoticity in Au+Au collisions at RHIC. Note that a similar study has recently been carried out for CERN LEP e^+e^- collisions [6] which reports a fully chaotic source.

The present three-pion correlation study by the STAR experiment at RHIC supplements the published two-pion correlation data from Au+Au collisions at $\sqrt{s_{NN}} = 130$ GeV [7]. A summary of the three-pion results will be presented for two multiplicity classes. By looking at collision classes with different multiplicities we can vary the impact parameter, and thus the number of initially colliding nucleons, and study the effect of the size of the colliding system on the source chaoticity. We discuss the method of normalization of the correlation function and its extrapolation to vanishing relative momentum in order to extract the source chaoticity; we estimate the various systematic uncertainties associated with these procedures. In principle, the relative momentum dependence of the three-pion correlator is sensitive to source asymmetries [3, 8], and we discuss the prospects for extracting the latter from our data.

Before presenting our experimental results, we first outline the formalism which guided our analysis (for details see Ref. [3] and references therein). The measured observable is the normalized three-pion correlator:

$$r_3\left(Q_3\right) = \frac{\left(C_3\left(Q_3\right)-1\right)-\left(C_2\left(Q_{12}\right)-1\right)-\left(C_2\left(Q_{23}\right)-1\right)-\left(C_2\left(Q_{31}\right)-1\right)}{\sqrt{\left(C_2\left(Q_{12}\right)-1\right)\left(C_2\left(Q_{23}\right)-1\right)\left(C_2\left(Q_{31}\right)-1\right)}} \tag{1}$$

Here $Q_3 = \sqrt{Q_{12}^2 + Q_{23}^2 + Q_{31}^2}$ and $Q_{ij} = \sqrt{-(p_i - p_j)^2}$ are the standard invariant relative momenta [4, 5] which can be computed for each pion triplet from the three measured momenta $(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3)$. $C_2\left(p_i, p_j\right) = \frac{P_2(p_i,p_j)}{P_1(p_i)P_1(p_j)} = C_2(Q_{ij})$ and $C_3\left(p_1,p_2,p_3\right) = \frac{P_3(p_1,p_2,p_3)}{P_1(p_1)P_1(p_2)P_1(p_3)} = C_3(Q_3)$, where P represents the momentum probability distribution. In Ref. [3] the ratio r_3 is defined in terms of functions which depend on all 9 components of $(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3)$; however, limited statistics even in our high-statistics sample requires a projection of both the numerator and denominator onto a single momentum variable, Q_3. We comment further on the implications of this projection below.

For fully chaotic sources $r_3/2$ approaches unity as all relative momenta (and thus Q_3) go to zero. If the source is partially coherent, a relationship can be established [3] between the limiting value of the three-pion correlator at $Q_3 = 0$ and the chaotic fraction ε $(0 \le \varepsilon \le 1)$ of the single-particle spectrum:

$$\frac{1}{2}\, r_3\left(Q_3{=}0\right) = \sqrt{\varepsilon}\, \frac{3-2\varepsilon}{(2-\varepsilon)^{3/2}}. \tag{2}$$

ε gives an upper limit on the value of the two-pion λ-parameter, which is sensitive to the fraction of coherent pairs in a sample, i.e. $\lambda = \varepsilon(2-\varepsilon)$ assuming no other effects on λ such as long-lived resonances[2]. Eq. (2) is not affected by the projection onto a single relative momentum variable. To exploit it and extract the degree of chaoticity ε, the measured data for r_3 must, however, be extrapolated from finite Q_3 to $Q_3 = 0$.

Similar to the two-boson correlation function, the three-boson correlation function $C_3\left(Q_3\right)$ is calculated from the data by taking the ratio $\frac{A(Q_3)}{B(Q_3)}$ and normalizing it to unity at large Q_3. Here $A\left(Q_3\right) = \frac{dN}{dQ_3}$ is the three-pion distribution as a function of the invariant three-pion relative momentum, integrated over the total momentum of the pion triplet as well as all other relative momentum components. It is obtained by taking three pions from a single event, calculating Q_3, and binning the results in a histogram. $B\left(Q_3\right)$ is the analogous mixed event distribution which is computed by taking a single pion from each of three separate events. Because of the zero in the denominator of the normalized three-pion correlator r_3 at large Q_{ij}, the particular method of normalization of C_2 and C_3 can have a huge effect on the calculation. The propagation of statistical errors through the r_3 functions, however, accounts for these effects completely. In fact, it is only with the enormous statistics available from STAR that the calculation can be considered in the range $15 < Q_3 < 120\,\mathrm{MeV}/c$. This range is large enough to provide reliable extrapolation to $Q_3 = 0$.

Data for the present results are from about 300K events taken during the $\sqrt{s_{NN}} = 130$ GeV Au+Au run at STAR using the Time Projection Chamber (TPC) [9] as the primary tracking detector. In the discussion that follows, all phase space cuts and experimental corrections are similar to the two-pion HBT analysis [7]. A set of multiplicity classes was created by taking the 12% most central for the high-multiplicity class and the next 20% most central for the mid-multiplicity class. For both multiplicity bins, tracks were constrained to have p_T in the range $0.125 < p_T < 0.5\,\mathrm{GeV}/c$, and pseudorapidity $|\eta| < 1.0$. A vertex cut was also applied to events such that the vertex along the z-axis (beam direction) had to fall within ± 75 cm of the center of the detector. In the range $15 < Q_3 < 120\,\mathrm{MeV}/c$, approximately 150 million triplets were included in both the negative and positive pion studies.

The C_2 correlation function was corrected for Coulomb repulsion with a finite Gaussian source approximation, using an integration of Coulomb wave functions [10]. This Coulomb effect was calculated for each mixed-event (only) pair. By thus weighting the mixed event distribution appropriately, the Coulomb effect is canceled in the ratio of real to mixed event distributions. In calculating C_3, the correction was applied by taking the product of three two-pion corrections, obtained from the three possible pairs formed from each mixed-event triplet. This type of correction approximates the three-body Coulomb problem to first order [11, 12]. A source size of 5 fm was used for both multiplicity bins, as was done in the two-pion analysis [7]. Other methods to more accurately estimate the true three-body Coulomb effect show a 5-10% smaller correction for a source radius of 5 fm [13]. This difference was applied to the Coulomb correction factor in calculating C_3, and the resulting shifts in the r_3 function were found to be within systematic uncertainties (these are shown in plots later in the paper). A separate study examined the effect of inappropriately applying the Coulomb correction to pions which come from long-lived resonances [14]. Using a rescattering model [15], the value of r_3 was found to increase by 10% when pairs and triplets of pions which contain pions from long-lived resonances were inappropriately Coulomb corrected. This is not included in the systematic uncertainty but will be taken into account later in the paper in the extraction of the chaotic fraction from r_3. Effects of finite momentum resolution on r_3 were also studied using this model and were found to be insignificant. The $1-\sigma$ uncertainty in determining Q_3 is found to be about 10 MeV/c.

Effects of track splitting during reconstruction were eliminated through a topological cut based on the TPC pad row configuration of hits composing a track, and track merging was corrected with a requirement on the minimum separation of a pair of tracks entering the TPC ($\geq$50 cm from the vertex). For C_3, for any pair within the triplet, if the pair entrance separation was less than 2.5 cm, the triplet was rejected. For the calculation of r_3, it was found that there was no significant deviation as the entrance separation was increased from 2.5 cm to 5.0 cm.



FIG. 1: Three-pion correlation function for central Au-Au events using π^- triplets. Statistical and statistical+systematic errors are shown.

Figure 1 shows the C_3 correlation function for negatively charged pions in the high-multiplicity bin. The shape of C_3 is mostly built up of products of two-pion correlations with the effect of true three-pion correlations being more subtle. At large Q_3, C_3 approaches unity and for an ideal pion source, i.e. $\lambda = 1$, C_3 would approach 6 at $Q_3 = 0$ (this is not the present case since $\lambda < 1$). A Gaussian parameterization is inadequate to describe this correlation function; this is consistent with results obtained in other experiments and a simulation[4, 5, 15]. In calculating r_3, the actual binned values of the correlation function for the various values of Q_3 are used instead of a fit [15]. In order to use Eq.(1), triplets are obtained that pass all of the momentum space and experimental cuts. Q_3, Q_{12}, Q_{23} and Q_{31} are calculated from the triplet and the three pairs that can be formed from the triplet. The values $C_3\,(Q_3)$, $C_2\,(Q_{12})$, $C_2\,(Q_{23})$ and $C_2\,(Q_{31})$ are then computed from the binned two- and three-pion correlation functions. These values are then used to calculate r_3, which is then binned as a function of Q_3. The average for each bin is then calculated to obtain the final result. Systematic uncertainties are greatest at the low Q_3 end due to track merging effects and the uncertainty in the Coulomb correction. The parameters controlling these effects were modified $\pm 20\%$ from the nominal values to obtain the overall systematic uncertainty in each bin.

The results for the two multiplicity bins are shown in Figure 2 for π^- and π^+, plotted as functions of Q_3^2. Plotting in this way is suggested by the theoretical analysis in [3] which shows that the leading relative momentum dependencies in the numerator and denominator of Eq. (1) are quadratic [16], allowing for a linear extrapolation of the results shown in Figure 2 to $Q_3 = 0$ by fitting them



FIG. 2: Calculation of r_3 for (a) central and (b) mid-central π^- events, and (c) central and (d) mid-central π^+ events. The fits shown use Eq. (3) to determine the intercept. Statistical and statistical+systematic uncertainties are shown.

to the form

$$r_3\left(Q_3\right)/2 = r_3(0)/2 - \alpha\, Q_3^2. \qquad (3)$$

where $r_3(0)/2$ and α are fit parameters. From Figure 2 it appears that the normalized three-pion correlator $r_3(Q_3)$ does indeed show a leading quadratic dependence for the smaller Q_3^2 values (Eq. (3) was fit to the range $0 < Q_3 < 60$ MeV/c).

The resulting intercepts $r_3(0)/2$ are shown in Figure 3, along with the results of WA98 and NA44. Error bars for STAR points are statistical+systematic. Intercepts from the quadratic fits as well as from quartic fits (i.e. adding a quartic term to Eq. (3) and fitting over the broader range $0 < Q_3 < 120$ MeV/c) are shown for comparison, and are seen to agree within errors. The STAR π^+ and π^- results are also seen to agree within error bars. NA44 reported a result close to unity for Pb-Pb interactions, but a much lower result for S-Pb [4], both with no clear Q_3 dependence. The Pb-Pb result from WA98 is somewhat smaller than that from NA44, although they agree within error bars, and the Q_3-dependence in their result is similar to what is seen in STAR [5]. A qualitatively similar, but much weaker relative momentum dependence of r_3 has been shown to exist in calculations by Nakamura and Seki for various parametrizations of chaotic sources [8]. In their model the relative momentum dependence is of higher than quadratic order, and it is an indication for an azimuthally asymmetric source. A direct comparison of our data with these model calculations is not possible due to our need for projecting the correlation functions on a single relative momentum variable. This introduces a leading quadratic dependence of r_3 on Q_3 [16] which buries the expected higher order effects arising from possible asymmetries of the emission function.

Figure 4 shows the results from the calculation of ε for STAR's measurements, and for those from WA98 and NA44, plotted versus charged particle multiplicity. The calculation was done starting with the results of Figure 3, decreasing them by 10% to approximately take into account the overcorrection produced by Coulomb-correcting long-lived resonances (see earlier discussion) and using Eq. (2). It was assumed that the 10% correction also applies to the SPS data, and to be conservative, a $\pm 5\%$ systematic uncertainty on the correction (i.e. $10\% \pm 5\%$) was included in all of the error bars shown. The plot shows an increasing trend in the STAR π^- and π^+ results going from mid-central to central. For the mid-central data, the results for ε show a partially chaotic source, as seen in the SPS results. The central data appear to give a mostly chaotic pion source. Including the SPS measurements into the overall systematics, there appears to be, within the uncertainties shown, a systematic increase in ε with increasing particle multiplicity, the smallest value being for SPS S-Pb collisions and the largest value for STAR central Au-Au collisions ($dN/d\eta$ for charged particles at mid-η for SPS S-Pb, SPS Pb-Pb, STAR mid-central, and STAR central are approximately 100, 370, 280, and 510, respectively[17, 18]). It is also found for the STAR results that the upper limit on the two-pion λ-parameter obtained from ε using the relationship mentioned earlier is in the range $0.71 - 0.81$ for mid-central and $0.91 - 0.97$ for central events. The actual values for λ extracted from STAR two-pion HBT measurements are 0.53 for mid-central and 0.50 for central events[7]. The lower λ-values extracted from the two-pion experiment can be explained in terms of long-lived resonance effects, which nicely cancel out in a three-pion analysis[15].

In summary, we have presented three-pion HBT results for $\sqrt{s_{NN}} = 130\,\mathrm{GeV}$ data at STAR, and have shown that for the central multiplicity class the STAR data indicate a large degree of chaoticity in the source at freeze-out, whereas for the mid-central class the source is less chaotic. Our r_3 results are close to those extracted in SPS Pb+Pb collisions, but differ from the low value obtained in SPS S+Pb collisions. The comparison between SPS and STAR results suggests a systematic dependence of the chaoticity on particle multiplicity. High statistics from STAR have allowed a normalized three-pion correlator calculation that extends to 120 MeV/c in Q_3, and the dependence on this variable has been shown to be quadratic in nature for low Q_3. STAR's measured values provide increased confidence in the validity of standard HBT analyses based on the assumption of a chaotic source for central collisions at RHIC.

We wish to thank the RHIC Operations Group and the RHIC Computing Facility at Brookhaven National Labo-



FIG. 3: Asymptotic value of $r_3/2$ from STAR and two other experiments [4, 5]. For STAR, (a) central and (b) mid-central results are shown for π^- (circular markers) and π^+ (square markers) data. The other experiments use π^- data only. STAR results for fitting with both a quadratic and quartic functions are shown.



FIG. 4: Chaotic fraction, calculated from Eq. (2), and plotted versus charged particle multiplicity for the same experiments as in Figure 3. The meanings of the symbols used in this figure are the same as in Figure 3. All results from Figure 3 are reduced by 10% in the calculations to approximately remove Coulomb-correction of pions from long-lived resonances.

ratory, and the National Energy Research Scientific Computing Center at Lawrence Berkeley National Laboratory for their support. This work was supported by the Division of Nuclear Physics and the Division of High Energy Physics of the Office of Science of the U.S. Department of Energy, the United States National Science Foundation, the Bundesministerium fuer Bildung und Forschung of Germany, the Institut National de la Physique Nucleaire et de la Physique des Particules of France, the United Kingdom Engineering and Physical Sciences Research Council, Fundacao de Amparo a Pesquisa do Estado de Sao Paulo, Brazil, the Russian Ministry of Science and Technology, the Ministry of Education of China, the National Natural Science Foundation of China, the Swiss National Science Foundation, and the Grant Agency of the Czech Republic.

---

[1] M. Gyulassy, S.K. Kauffmann, and L.W. Wilson, Phys. Rev. C **20**, 2267 (1979).
[2] U. A. Wiedemann and U. Heinz, Phys. Rep. **319**, 145 (1999).
[3] U. Heinz and Q. H. Zhang, Phys. Rev. C **56**, 426 (1997).
[4] I.G. Bearden *et. al.*, Phys. Lett. B **517**, 25 (2001).
[5] M.M. Aggarwal *et. al.*, Phys. Rev. Lett. **85**, 2895 (2000).
[6] P. Achard *et. al.*, Phys. Lett. B **540**, 185 (2002).
[7] C. Adler *et. al.*, Phys. Rev. Lett. **87**, 082301 (2001).
[8] H. Nakamura and R. Seki, Phys. Rev. C **60**, 064904 (1999).
[9] K. H. Ackermann *et al.*, Nucl. Instrum. Meth. A **499**, 624 (2003).
[10] S. Pratt, T. Csörgő and J. Zimányi, Phys. Rev. C **42**, 2646 (1990).
[11] S.P. Merkuriev, Theor. Math. Phys. **32**, 680 (1977).
[12] M. Brauner, J.S. Briggs and H. Klar, J. Phys. B **22**, 2265 (1989).
[13] E.O. Alt, T. Csörgő, B. Lörstad, J. Schmidt-Sorensen, Phys. Lett. B **458**, 407 (1999).
[14] T. J. Humanic, preprint to be submitted.
[15] T.J. Humanic, Phys. Rev. C **60**, 014901 (1999).
[16] For a fully chaotic source ($\varepsilon = 1$), the leading quadratic relative momentum dependencies in the numerator and denominator of Eq. (1) cancel, if the ratio is taken pointwise in 9-dimensional $(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3)$ space. If numerator and denominator are first projected on the single variable Q_3, this cancellation in the ratio is spoiled. For partially coherent sources there is no such cancellation to begin with [3]. The parametrization Eq. (3) thus captures the expected leading Q_3 dependence of the data (U. Heinz, private communication).
[17] H. Appelshäuser *et. al.*, Phys. Rev. Lett. **82**, 2471 (1999).
[18] B. B. Back *et. al.*, Phys. Rev. C **65**, 061901 (2002).