Interplay of correlations & fluctuations at RHIC — What do we learn?

Mikhail Kopytine (Kent State University)

for the STAR Collaboration

Argonne National Laboratory – USA, Brookhaven National Laboratory – USA, U. of Birmingham – UK, U. of California, Berkeley – USA, U. of California, Davis – USA, U. of California, Los Angeles – USA, Carnegie Mellon U. – USA, Creighton U. – USA, Laboratory for High Energy, JINR – Russia, Particle Physics Laboratory, JINR – Russia, U. of Frankfurt – Germany, Indiana U. – USA, Institute de Recherches Subatomiques – France, Kent State U. – USA, Lawrence Berkeley National Laboratory – USA, Max-Plank-Institute für Physik – Germany, Michigan State U. – USA, Moscow Engineering Physics Institute – Russia, City College of New York – USA, Ohio State U. – USA, Pennsylvania State U. – USA, Institute of High Energy Physics – Russia, Purdue U. – USA, Rice U. – USA, Universidade de Sao Paolo – Brazil, SUBATECH – France, Texas A & M – USA, U. of Texas – USA, Warsaw U. of Technology – Poland, U. of Washington – USA, Wayne State U. – USA, Institute of Particle Physics – China, and Yale U. – USA

Correlations & fluctuations... how to learn anything?

- Discrete Wavelet Transform $\Rightarrow$ power spectra of local track density fluctuations
- 'thermal' reference = mixed events
- How does the medium change with initial conditions ? In this language:
  - thermalization: reality $\rightarrow$ reference
  - correlation: reality $<$ reference
  - fluctuation: reality $>$ reference (e.g. jets)

Mikhail Kopytine *Kent State University*





$F^{\lambda}_{m,i,j}(\phi,\eta)$–Haar wavelet basis in 2D:



scale fineness (m), directional modes of sensitivity (λ), track density $\rho(\eta,\phi,p_T)$.

Basic observables:

Power of local fluctuations, mode λ:

$$P^{\lambda}(m) = \frac{1}{2^{2m}} \sum_{i,j} \langle \rho, F^{\lambda}_{m,i,j} \rangle^2, \tag{1}$$

Dynamic texture:

$$P^{\lambda}(m)_{true} - P^{\lambda}(m)_{mix} \tag{2}$$

"incoherently" normalized :

$$(P^{\lambda}(m)_{true} - P^{\lambda}(m)_{mix})/P^{\lambda}(m)_{mix} \tag{3}$$

"coherently" normalized :

$$(P^{\lambda}(m)_{true} - P^{\lambda}(m)_{mix})/P^{\lambda}(m)_{mix}/N, \tag{4}$$

where N is (sub)event (p_T bin) multiplicity

Scale is localized:

a)





b)





c)







Event mixing scheme:

– no pixel is used twice

– not more than one pixel from any given true event per mixed event

– no mixing of "different" events: multiplicity, vertex

With n pixels, need $\geq n^2$ events per event class.



Proper **normalization** $\Rightarrow$ results independent of bin size or subevent multiplicity.

Coherent: $(P^{\lambda}(m)_{true} - P^{\lambda}(m)_{mix})/P^{\lambda}(m)_{mix}/N$,

Incoherent: $(P^{\lambda}(m)_{true} - P^{\lambda}(m)_{mix})/P^{\lambda}(m)_{mix}$



HIJING – azimuthal back to back correlations (fineness=2 is enhanced!) ● ○– scale 1; ● ○ – scale 2.





● ○– scale 1; ● ○ – scale 2. HIJING shows coherent scaling with p_T bin width



$\sqrt{S_{NN}}$ = 130 GeV. Fineness scale=1 ($\delta\eta$ = 1); **Charged hadrons:** ⋆ – STAR; • – regular HIJING; ∘ – HIJING without jets. The HIJING texture is mostly jets. STAR data: a change of regime with centrality.



$\sqrt{S_{NN}} = 130$ GeV. Fineness scale=1 ($\delta\eta = 1$);
$P_{true} < P_{mix} \Rightarrow$ long range correlation



What is the origin of the long range η correlation = suppressed fluctuation ? Need common memory/communication over $\delta\eta \approx 1$. Different break-up points **causally disjoint.** Hadron rescattering can not increase order.



Fineness scale=1 ($\delta\eta = 1$). Peripheral ($mult/n_0 < 0.1$) events: $\star$ – STAR data for $\sqrt{S_{NN}} = 200$ GeV. $\bullet$ – HIJING @ same energy.



Fineness scale=1 ($\delta\eta = 1$). $\star$ – STAR $\sqrt{S_{NN}} =$ 200 GeV; $\bullet$ – regular HIJING; $\circ$ – HIJING+jet quenching (both 130 GeV). Central ($0.65 < mult/n_0 < 1$) events: Data more 'thermal' at $p_T > 0.6 GeV \rightarrow$ dissipation?

Bose-Einstein/Coulomb contribution at low p_T needs to be quantified.

Conclusions:

– peripheral ($mult/n_0 < 0.1$) data qualitatively agree with HIJING except for elliptic flow; details of p_T behaviour differ

– long range correlation (suppressed fluctuation) in η becomes visible in central events.

– p_T-dependence of η-texture: at soft $0.6 < p_T < 2$ GeV, jet texture is suppressed. At low p_T, need to understand the "HBT" contribution.

– more info: **nucl-ex/0211015, nucl-ex/0211019**