




RHIC and the road to an EIC



Matthew A. C. Lamont
Brookhaven National Lab

Most compelling physics questions


Spin physics



- What is the polarisation of gluons at small x where they dominate?
- What is the x -dependence and flavour decomposition of the polarised sea?


Determine quark and gluon contributions to the proton spin at last!!

Imaging




- What is the spatial distribution of quarks/ gluons in nucleons AND nuclei?
- Understand deep aspects of gauge theories revealed by k_T dependent distributions

Possible window to orbital angular momentum




Strong Colour Fields and Hadronisation



- Quantitatively probe the universality of strong colour fields in $A+A$, $p+A$ and $e+A$
- Understand in detail the transition to the non-linear regime of strong gluon fields and the physics of saturation
- How do hard probes in $e+A$ interact with the medium?



Currently have no experimental knowledge of gluons in nuclei at small x !!

Extension of x, Q^2 coverage with an EIC



- Increase reach in x by a factor of 100 in both polarised $e+p$ and $e+A$ - into the range where gluons dominate
 - ➔ $e+p$: constrain the helicity sum rules?
 - ➔ $e+A$: saturation effects become visible?
- Increase in Q^2 coverage
 - ➔ study scaling violations


Extension of x, Q^2 coverage with an EIC



- Increase reach in x by a factor of 100 in both polarised e+p and e+A - into the range where gluons dominate
 - ➔ e+p: constrain the helicity sum rules?
 - ➔ e+A: saturation effects become visible?
- Increase in Q^2 coverage
 - ➔ study scaling violations




Constraining $\Delta g(x)$ at RHIC, EIC




- RHIC data can constrain $\Delta g(x)$ down to a few $\times 10^{-2}$
 - ➔ Latest RHIC data show non-zero $\Delta g(x)$ in measured range
 - ➔ Large unmeasured region still exists

Constraining $\Delta g(x)$ at RHIC, EIC



- RHIC data can constrain $\Delta g(x)$ down to a few $x \cdot 10^{-2}$
 - ➔ Latest RHIC data show non-zero $\Delta g(x)$ in measured range
 - ➔ Large unmeasured region still exists
- DIS measurements at an EIC will lead to a dramatic reduction of the uncertainties in the unmeasured region

Constraining $\Delta g(x)$ at RHIC, EIC



- RHIC data can constrain $\Delta g(x)$ down to a few $\times 10^{-2}$
 - ➔ Latest RHIC data show non-zero $\Delta g(x)$ in measured range
 - ➔ Large unmeasured region still exists
- DIS measurements at an EIC will lead to a dramatic reduction of the uncertainties in the unmeasured region



Nuclear DIS → Structure Functions

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2) \quad x = \frac{Q^2}{2pq} = \frac{Q^2}{sy}$$

Strategies:

slope of y^2/Y_+ for different s at fixed x & Q^2

e+Au: 1st stage

5x50 - A/Ldt = 2 fb⁻¹

5x75 - A/Ldt = 4 fb⁻¹

5x100 - A/Ldt = 4 fb⁻¹


running combined

~6 months total running

(50% eff)

statistical errors are swamped by
the 1% systematic errors

Will be dominated by systematics,
but would need a full detector
simulation in order to estimate them



Region with non-linear effects
should be seen at EIC!!



Nuclear DIS → Structure Functions

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2) \quad x = \frac{Q^2}{2pq} = \frac{Q^2}{sy}$$

Strategies:

slope of y^2/Y_+ for different s at fixed x & Q^2

e+Au: 1st stage

5x50 - $A\mathcal{L}dt = 2 \text{ fb}^{-1}$

5x75 - $A\mathcal{L}dt = 4 \text{ fb}^{-1}$

5x100 - $A\mathcal{L}dt = 4 \text{ fb}^{-1}$





running combined

~6 months total running

(50% eff)

statistical errors are swamped by the 1% systematic errors

Will be dominated by systematics, but would need a full detector simulation in order to estimate them





Nuclear DIS → Structure Functions

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2) \quad x = \frac{Q^2}{2pq} = \frac{Q^2}{sy}$$

Strategies:

slope of y^2/Y_+ for different s at fixed x & Q^2

e+Au: 1st stage

5x50 - $A \int L dt = 2 \text{ fb}^{-1}$

5x75 - $A \int L dt = 4 \text{ fb}^{-1}$


5x100 - $A \int L dt = 4 \text{ fb}^{-1}$

running combined

~6 months total running
(50% eff)


statistical errors are swamped by
the 1% systematic errors

Will be dominated by systematics,
but would need a full detector
simulation in order to estimate them




SIDIS in e+p → flavour-separated helicity PDFs

- SIDIS measurements with identified π , k lead to much reduced uncertainties in the flavour-separated helicity PDFs as in $\Delta g(x)$







SIDIS in $e+A \rightarrow$ di-hadron correlations





- Long history of di-hadron correlation measurements in p+p, d+A and A+A collisions
 - ➔ central rapidity show attenuation of away-side correlation in A+A but not p+p nor d+A collisions
 - Final state effect

SIDIS in $e+A \rightarrow$ di-hadron correlations



- Long history of di-hadron correlation measurements in p+p, d+A and A+A collisions
 - ➔ central rapidity show attenuation of away-side correlation in A+A but not p+p nor d+A collisions
 - Final state effect
 - ➔ forward di-hadron correlations show an attenuation even in d+Au
 - Best evidence so far for manifestation of saturation effects

SIDIS in $e+A \rightarrow$ di-hadron correlations




We don't know the x , Q^2 in d+A!!

- Long history of di-hadron correlation measurements in p+p, d+A and A+A collisions
 - ➔ central rapidity show attenuation of away-side correlation in A+A but not p+p nor d+A collisions
 - Final state effect
 - ➔ forward di-hadron correlations show an attenuation even in d+Au
 - Best evidence so far for manifestation of saturation effects



SIDIS in $e+A \rightarrow$ di-hadron correlations




Xiao, Yuang et al (private communication)

- Predictions from a saturation model show an ordered attenuation of the away-side with increasing nuclear mass
 - Simulations (PYTHIA + DPMJETIII) for e+Au show that the sat/no-sat scenarios can be distinguished within errors
- ➡ Gives a handle on multi-gluon distributions




SIDIS in $e+A \rightarrow$ di-hadron correlations



Xiao, Yuang et al (private communication)


- Predictions from a saturation model show an ordered attenuation of the away-side with increasing nuclear mass
 - Simulations (PYTHIA + DPMJETIII) for $e+Au$ show that the sat/no-sat scenarios can be distinguished within errors
- ➡ Gives a handle on multi-gluon distributions

Exclusive processes in e+A - diffraction



- β is the momentum fraction of the struck parton w.r.t. the Pomeron
- $x_{IP} = x/\beta$: momentum fraction of the exchanged object (Pomeron) w.r.t. the hadron

$$\beta = \frac{x}{x_{IP}} = \frac{Q^2}{Q^2 + M_X^2 - t}$$





• Diffraction in e+p:

- ➔ HERA: 15% of all events are diffractive

• Diffraction in e+A:


- ➔ Predictions: $\sigma_{\text{diff}}/\sigma_{\text{tot}}$ in e+A ~25-40%
- ➔ Coherent diffraction (nuclei intact)
- ➔ Incoherent diffraction: breakup into nucleons (nucleons intact)

Day 1: Diffractive Cross-sections




- Ratio of diffractive-to-total cross-section drastically different between saturation (Marquet) and non-saturation (Frankfurt, Guzey, Strikman) models
- Expected experimental error bars (simulated for 10 fb^{-1} of data) can distinguish between the two scenarios

Exclusive vector meson production




$$d\sigma \propto g(x)^2$$




- Exclusive vector meson production is most sensitive to the gluon distribution
 - ➔ colour-neutral exchange of gluons
- J/ψ shows some difference between saturation and no-saturation
- ϕ shows a much larger difference
 - ➔ wave function for ϕ is larger and hence more sensitive to saturation effects

Exclusive vector meson production





$$d\sigma \propto g(x)^2$$



- Exclusive vector meson production is most sensitive to the gluon distribution
 - ➔ colour-neutral exchange of gluons
- J/ψ shows some difference between saturation and no-saturation
- ϕ shows a much larger difference
 - ➔ wave function for ϕ is larger and hence more sensitive to saturation effects

Exclusive Vector Meson Production in e+A



- Low-t: coherent diffraction dominates - **gluon density**
- High-t: incoherent diffraction dominates - **gluon correlations**
- ➔ Need good breakup detection efficiency to discriminate between the two scenarios
 - unlike protons, forward spectrometer won't work for heavy ions
 - measure emitted neutrons in a ZDC
 - rapidity gap with absence of break-up fragments sufficient to identify coherent events





Finding the source...

- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)







Finding the source...

- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)





Finding the source...

- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)





Finding the source...

- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)





Finding the source...

- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)





Finding the source...

- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)





Finding the source...

- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)





Finding the source...


- Take the $d\sigma/dt$ distribution and perform a Fourier Transform to extract the b -distribution of the gluons

$$F(b) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$ (for small x)



Imaging in $e+p$ - DVCS





Summary/Conclusions

BACKUP




The pillars of the eRHIC physics programme



- Wide physics programme with demanding requirements on detector and machine performance




The pillars of the eRHIC physics programme



- Wide physics programme with demanding requirements on detector and machine performance



The pillars of the eRHIC physics programme



- Wide physics programme with demanding requirements on detector and machine performance




Outline

- Why an EIC?
 - ➔ Spin + e+A
- What is eRHIC?
- What can eRHIC do for you?
- How do we realise eRHIC?

The eRHIC project

- eRHIC:


- ➔ Utilises the RHIC ion beams
- ➔ Two 2.45 GeV Energy Recovery Linacs (ERLs) accelerate the e^- beam
 - ▶ 6 separate rings accelerate the e^- up to a maximum energy of 30 GeV
- ➔ 2-stage approach
 - ▶ Stage 1: e^- 5-10 GeV
 - ▶ Stage 2: e^- 20-30 GeV
- ➔ Space for new detector at IP12
 - ▶ Possibilities for collisions in current STAR and PHENIX IPs



The eRHIC project

- eRHIC:


- ➔ Utilises the RHIC ion beams
- ➔ Two 2.45 GeV Energy Recovery Linacs (ERLs) accelerate the e^- beam
 - ▶ 6 separate rings accelerate the e^- up to a maximum energy of 30 GeV
- ➔ 2-stage approach
 - ▶ Stage 1: e^- 5-10 GeV
 - ▶ Stage 2: e^- 20-30 GeV
- ➔ Space for new detector at IP12
 - ▶ Possibilities for collisions in current STAR and PHENIX IPs



The eRHIC project

- eRHIC:


- ➔ Utilises the RHIC ion beams
- ➔ Two 2.45 GeV Energy Recovery Linacs (ERLs) accelerate the e^- beam
 - ▶ 6 separate rings accelerate the e^- up to a maximum energy of 30 GeV
- ➔ 2-stage approach
 - ▶ Stage 1: e^- 5-10 GeV
 - ▶ Stage 2: e^- 20-30 GeV
- ➔ Space for new detector at IP12
 - ▶ Possibilities for collisions in current STAR and PHENIX IPs



The eRHIC project

- eRHIC:


- ➔ Utilises the RHIC ion beams
- ➔ Two 2.45 GeV Energy Recovery Linacs (ERLs) accelerate the e^- beam
 - ▶ 6 separate rings accelerate the e^- up to a maximum energy of 30 GeV
- ➔ Space for new detector at IP12
 - ▶ Possibilities for collisions in current STAR and PHENIX IPs



The eRHIC project

- eRHIC:


- ➔ Utilises the RHIC ion beams
- ➔ Two 2.45 GeV Energy Recovery Linacs (ERLs) accelerate the e^- beam
 - ▶ 6 separate rings accelerate the e^- up to a maximum energy of 30 GeV
- ➔ Space for new detector at IP12
 - ▶ Possibilities for collisions in current STAR and PHENIX IPs



The eRHIC project

- eRHIC:


- ➔ Utilises the RHIC ion beams
- ➔ Two 2.45 GeV Energy Recovery Linacs (ERLs) accelerate the e^- beam
 - ▶ 6 separate rings accelerate the e^- up to a maximum energy of 30 GeV
- ➔ Space for new detector at IP12
 - ▶ Possibilities for collisions in current STAR and PHENIX IPs



The eRHIC project


- eRHIC:

- ➔ Utilises the RHIC ion beams
- ➔ Two 2.45 GeV Energy Recovery Linacs (ERLs) accelerate the e^- beam
 - ▶ 6 separate rings accelerate the e^- up to a maximum energy of 30 GeV
- ➔ Space for new detector at IP12
 - ▶ Possibilities for collisions in current STAR and PHENIX IPs











(Very) Brief Recap of Saturation at an EIC




(Very) Brief Recap of Saturation at an EIC






(Very) Brief Recap of Saturation at an EIC



$$Q_s^2(x) \sim A^{1/3} \left(\frac{1}{x} \right)^\lambda$$



(Very) Brief Recap of Saturation at an EIC



$$Q_s^2(x) \sim A^{1/3} \left(\frac{1}{x} \right)^\lambda$$

